

■ CK20 weak (%) ■ CK20 mod. (%) ■ CK20 str. (%)

% of CK7 negative tumor samples

0.0 10.0 20.0 30.0 40.0 50.0 60.0 70.0 80.0 90.0 100.0

Adenomatous polyp, low-grade dysplasia (n=42)



Non-invasive papillary urothelial carcinoma, pTa G2 low grade (n=2)

Adenomatous polyp, high-grade dysplasia (n=45)

Adenocarcinoma of the colon (n=1626)

Mucinous carcinoma of the ovary (n=6)

Non-invasive papillary urothelial carcinoma, pTa G3 (n=6)

Pancreatic/Ampullary adenocarcinoma (n=15)

Non-invasive papillary urothelial carcinoma, pTa G2 high grade (n=5)

Merkel cell carcinoma (n=37)

Gastric adenocarcinoma, diffuse type (n=27)

Adenocarcinoma of the esophagus (n=10)

Gastric adenocarcinoma, intestinal type (n=62)

Gastric adenocarcinoma, mixed type (n=12)

Acinar cell carcinoma of the pancreas (n=4)

Endometrioid carcinoma of the ovary (n=5)

Ductal adenocarcinoma of the pancreas (n=5)

Adenocarcinoma of the prostate, Gleason 4+4 (n=63)

Urothelial carcinoma, pT2-4 G3 (n=110)

Appendix, neuroendocrine tumor (NET) (n=15)

Colorectal, neuroendocrine carcinoma (NEC) (n=8)

Adenocarcinoma of the prostate, Gleason 3+3 (n=57)

Pancreas, neuroendocrine tumor (NET) (n=77)

Pancreas, neuroendocrine carcinoma (NEC) (n=11)

Adenocarcinoma of the prostate, Gleason 5+5 (n=77)

Teratoma (n=12)

Adenocarcinoma of the prostate (recurrence) (n=221)

Hepatocellular carcinoma (n=34)

Endometrioid endometrial carcinoma (n=22)

Ileum, neuroendocrine tumor (NET) (n=46)

Papillary renal cell carcinoma (n=35)

Squamous cell carcinoma of the esophagus (n=37)

Squamous cell carcinoma of the larynx (n=77)

Squamous cell carcinoma of the vagina (n=49)

Clear cell renal cell carcinoma (n=663)