
CK7 weak (%)
CK7 mod. (%)
CK7 str. (%)
% of CK20 positive tumor samples
0.0
10.0
20.0
30.0
40.0
50.0
60.0
70.0
80.0
90.0
100.0
Oral squamous cell carcinoma (floor of the mouth) (n=1)
Adenocarcinoma of the lung (n=9)
Carcinosarcoma of the uterus (n=1)
Serous carcinoma of the ovary (n=1)
Invasive breast carcinoma of no special type (n=7)
Medullary carcinoma of the breast (n=1)
Squamous cell carcinoma of the anal canal (n=2)
Cholangiocarcinoma (n=10)
Sarcomatoid urothelial carcinoma (n=1)
Ductal adenocarcinoma of the pancreas (n=113)
Non-invasive papillary urothelial carcinoma, pTa G2 low grade (n=105)
Non-invasive papillary urothelial carcinoma, pTa G2 high grade (n=98)
Urothelial carcinoma, pT2-4 G3 (n=398)
Non-invasive papillary urothelial carcinoma, pTa G3 (n=110)
Mucinous carcinoma of the ovary (n=52)
Endometrioid endometrial carcinoma (n=8)
Endometrioid carcinoma of the ovary (n=8)
Adenocarcinoma of the esophagus (n=32)
Gastric adenocarcinoma, mixed type (n=16)
Pancreatic/Ampullary adenocarcinoma (n=38)
Squamous cell carcinoma of the vagina (n=3)
Gastric adenocarcinoma, diffuse type (n=57)
Papillary renal cell carcinoma (n=3)
Hepatocellular carcinoma (n=4)
Teratoma (n=2)
Gastric adenocarcinoma, intestinal type (n=64)
Squamous cell carcinoma of the larynx (n=3)
Adenocarcinoma of the prostate, Gleason 4+4 (n=14)
Pancreas, neuroendocrine tumor (NET) (n=11)
Adenocarcinoma of the prostate (recurrence) (n=20)
Adenocarcinoma of the prostate, Gleason 3+3 (n=9)
Merkel cell carcinoma (n=31)
Adenomatous polyp, low-grade dysplasia (n=48)
Adenocarcinoma of the prostate, Gleason 5+5 (n=8)
Adenocarcinoma of the colon (n=1647)
Adenomatous polyp, high-grade dysplasia (n=47)